



New Mexico

Live births, abortions, and pregnancies


Live birth rates, 1982–1992



Live birth rates,* 1991



Pregnancy rates, 1990



National live birth rate ranking, 1990


Age group		
15-17	18-19	15-19
10th highest	Highest	4th highest

*Missing bar indicates fewer than 10,000 females or fewer than 20 births. This graph refers to females aged 15–19 years.




Sexually transmitted diseases New Mexico

Gonorrhea rates, 1982–1992



Gonorrhea rates, 1992



National ranking,* 1992


Males – 40th highest Females – 38th highest

*National ranking refers to males and females aged 10–24 years.




Chlamydia cases and rates,* 1988–1992


Cases




Rates




Cases



Rates



Acquired immunodeficiency syndrome (AIDS) cumulative cases,† 1981–1992



*Chlamydia cases and rates refer to males and females of all ages.


†Missing bar indicates fewer than six cumulative cases.



Related risk behaviors,* 1993

New Mexico

Had sexual intercourse (lifetime)




Had sexual intercourse with four or more people (lifetime)



Had sexual intercourse during the past three months (currently sexually active)



Used a condom during last sexual intercourse (among currently sexually active)



Used birth control pills during last sexual intercourse (among currently sexually active)



Injected illegal drugs (lifetime)



*Related risk behaviors refer to high school students.

Sources: CDC—NCHS and NCCDPHP (birth, abortion, and pregnancy data), NCPS (gonorrhea and chlamydia data), NCID (AIDS data); 1993 New Mexico Youth Risk Behavior Survey (risk behavior data).

